



SOUTH WEST ZONE SUMMARY SHEET

KEY FINDINGS

MARCH 2019

The 2018 Nigeria HIV/AIDS Indicator and Impact Survey (NAIIS) was a national household-based survey that assessed the prevalence of human immunodeficiency virus (HIV) and related health indicators. Data collection in South West Zone was done from July through December 2018. Data were collected from household members age 0-64 years. Home-based HIV counseling and testing services were provided to people who participated in the survey. Participants receiving an HIV-positive test result were linked to treatment services. NAIIS data includes national, zonal, and state information on HIV control activities in Nigeria.

NAIIS was led by the Government of Nigeria through the Federal Ministry of Health (FMoH) and the National Agency for the Control of AIDS (NACA), conducted with funding from the United States (U.S.) President’s Emergency Plan for AIDS Relief (PEPFAR) and the Global Fund to Fight AIDS, Tuberculosis and Malaria with technical assistance from the U.S. Centers for Disease Control and Prevention (CDC). The survey was implemented by the NAIIS Consortium, led by the University of Maryland, Baltimore (UMB) under the supervision of the NAIIS Technical Committee.

SURVEY RESULTS



HIV Indicator	Female		Male		Total		Unweighted sample size
	%	95%CI*	%	95%CI	%	95%CI	
HIV prevalence †							
15–49 years	1.4	1.2–1.7	0.7	0.5–0.9	1.1	0.9–1.9	22,248
15-64 years	1.5	1.3–1.8	0.8	0.6–1.0	1.2	1.0–1.4	26,867
Viral load suppression ‡							
15–49 years	49.7	41.8–57.5	25.9	13.6–38.3	43.0	35.8–50.3	242
15-64 years	49.2	42.2–56.2	29.4	18.9–39.9	43.1	36.7–49.5	325

* The 95% CI (confidence interval) indicates the interval within which the true population parameter is expected to fall 95% of the time.

† The numerator for HIV prevalence is the number of people tested HIV-positive in each subgroup. The denominator is the number of people tested in each subgroup.

‡ Viral load suppression is defined as HIV RNA <1,000 copies per ml of plasma. The denominator for viral suppression is the number of PLHIV in each age group.


Prevalence of HIV among adults age 15-64 years was 1.2%, 1.5% among females and 0.8% among males.

Prevalence of viral load suppression (VLS) among people living with HIV (PLHIV) age 15-64 years in South West Zone was 43.1%, 49.2% among females and 29.4% among males.

HIV PREVALENCE BY SEX AND AGE



HIV prevalence was the highest among females age 45-49 years at 2.6% and the highest among males age 45-49 years at 2.1%. The HIV prevalence gender disparity between females and males was greatest among younger adults, with females age 20-24 years (0.9%) having slightly over 4 times the prevalence of males in the same age group (0.2%).




* The error bars show the 95% CIs (confidence intervals), the intervals within which the true population parameter is expected to fall 95% of the time.

HIV PREVALENCE AMONG PERSONS AGE 15-64 YEARS BY ZONE AND STATE




Among adults age 15-64 years, HIV prevalence varied by state across South West Zone, ranging from 1.6% in Ogun State to 0.8% in Ekiti State.



	HIV Prevalence (%)	95% CI*
South West	1.2	1.0-1.4
Ogun	1.6	1.1-2.1
Lagos	1.4	1.1-1.7
Ondo	1.1	0.7-1.5
Osun	0.9	0.6-1.3
Oyo	0.9	0.6-1.3
Ekiti	0.8	0.5-1.1


* The 95% CI (confidence interval) indicates the interval within which the true population parameter is expected to fall 95% of the time.



VIRAL LOAD SUPPRESSION AMONG PLHIV AGE 15-64 YEARS BY SEX AND AGE



VLS among PLHIV was the highest among females age 45-54 years at 66.6% and the highest among males age 55-64 years at 47.7%.




* VLS (viral load suppression) is defined as HIV RNA <1,000 copies per ml of plasma. The denominator for viral suppression is the number PLHIV in each age group.
 † The error bars show the 95% CIs (confidence intervals), the intervals within which the true population parameter is expected to fall 95% of the time.
 ‡ The estimates for females age 15-24 years and 55-64 years and males age 15-24 years, 25-34 years, 35-44 years, and 55-64 years were not presented because the unweighted sample size was <30 people. The estimates for males age 45-54 years, were based on an unweighted sample size of 30-49 people and should be interpreted with caution.

VIRAL LOAD SUPPRESSION AMONG PLHIV AGE 15-64 YEARS BY ZONE AND STATE, NAIS 2018



Among PLHIV age 15-64 years, VLS varied by state across South West Zone, ranging from 44.9% in Lagos State to 37.3% in Osun State.



	Viral Load Suppression (%)†	95% CI‡
South West	43.1	36.7–49.5
Ekiti	NA§	
Lagos	44.9	34.6–55.2
Ogun	44.7	32.0–57.4
Oyo	43.7	24.7–62.7
Ondo	39	23.1–54.9
Osun	37.3	20.5–54.2

† Viral load suppression is defined as HIV RNA <1,000 copies per ml of plasma. The denominator for viral suppression is the number PLHIV age 15-64 years.
 ‡ The 95% CI (confidence interval) indicates the interval within which the true population parameter is expected to fall 95% of the time.
 § NA – Data not presented due to an unweighted sample size less than 30 people.
 ¶ Estimates in parentheses were based on an unweighted sample size of 30-49 people and should be interpreted with caution.



Legend
 Viral Load Suppression (%)

- 16.0-34.9
- 35.0-38.9
- 39.0-43.9
- 44.0-52.9
- 53.0-75.1
- NA*

* NA – Data not presented due to an unweighted sample size of less than 30 people.

CONCLUSIONS



- HIV prevalence estimates varied by state in South West Zone, with the highest prevalence in Ogun State and the lowest prevalence in Ekiti State.
- In South West Zone, slightly more than 4 out of 10 PLHIV achieved viral suppression.
- South West Zone will focus efforts on ensuring those infected with achieve viral suppression, minimizing the risk for HIV transmission and moving Nigeria closer to controlling the HIV epidemic.

RESPONSE RATES AND HIV TESTING METHODS



Of 17,621 eligible households in South West Zone, 93.5% completed the household interview. Of 19,032 eligible women and 16,058 eligible men age 15-64 years, 81.1% of women and 71.2% of men were both interviewed and tested for HIV.

HIV prevalence testing was conducted in each household using a serological rapid diagnostic testing algorithm based on Nigeria's National HIV Testing Guidelines, with laboratory confirmation of seropositive samples using a supplemental assay.

The Government of Nigeria is grateful to all citizens who agreed to be part of NAIIS. Their dedication and willingness will help improve the lives of all Nigerians.



NAIIS is supported by PEPFAR through CDC under the terms of cooperative agreement GH18-1813, GH002108 and by the Global Fund to Fight AIDS, Tuberculosis and Malaria under contract NGA-H-NACA. The findings in this report should be considered preliminary and are subject to change. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the funding agencies.